



# Trigger Finger Release

## Patient Postoperative Instructions

### Laith Al-Shihabi, MD

#### **0-2 weeks**

##### Pain medication:

- Use your prescription pain medication as needed to control pain.
- Begin taking pain medication before the local anesthetic around your surgery site wears off.
- Do not drive or operate machinery while taking narcotic/opioid pain medication.
- Unless you cannot take them for other medical reasons, it is ok to use over-the-counter anti-inflammatory medication (e.g. Advil or Aleve) in addition to your prescription pain medication. Do not take tylenol as your prescription already contains this medication.

##### Post-op Dressings:

- Keep the dressings in place for 3 days after surgery, then remove them and cover the incision with a waterproof band-aid.
- Elevation of the hand, along with ice on-and-off for 20 minutes at a time, will help with swelling and pain.
- It is ok to shower within the first 3 days after surgery, but the dressings must stay clean and dry. Place a waterproof covering over the dressings while bathing or showering (such as a heavy plastic bag that is sealed with Saran wrap). After removing the dressing, it is ok to let water run over the incision with a shower or when washing your hands, but do not immerse the incision (e.g. no swimming or bath tub) or scrub it.
- If the dressings get dirty, wet, or come apart please contact the office.

##### Motion:

- Start moving your fingers and the thumb as soon as possible.
- You can use your hand for light activity, such as eating, drinking, or using a computer.
- You do not have any lifting or carrying restrictions, but higher demand activities may cause surgical-site pain. Avoid applying strong pressure to your palm in the area of your incision. Perform activities as your symptoms allow.

#### **2-6 weeks (First post-op visit)**

##### Splint:

- Sutures, if present, will be removed at your 2-week visit.
- No splint is necessary.
- You can begin gentle scar massage at 3 weeks after surgery.

##### Therapy:

- No therapy is necessary unless you have difficulty recovering your range of motion.

#### **6 -10 weeks (Second post-op visit)**

Schedule a second follow-up visit at 6 weeks in case you are still having problems or other symptoms. If you are doing well, no additional follow-up is necessary.